

DESY 98-140

8 September 1998

Rev. 4 January 1999

Enhancing Squark/Leptoquark Production by Increasing the HERA Beam Energies

Michiel Botje

NIKHEF

Günter Wolf

Deutsches Elektronen Synchrotron DESY

Abstract

The cross section for production of R-parity violating squarks ($\tilde{q}$) or leptoquarks (LQ) with masses above 200 GeV by positron-proton scattering at HERA can be substantially increased by a moderate increase of the beam energies. If, for example, the positron energy is raised from 27.5 to 32.5 GeV and the proton energy from 820 to 1000 GeV, the $\tilde{q}/LQ$ production cross section for masses of 200 – 230 GeV increases by a factor of about 2–6 and the search range for $\tilde{q}/LQ$ is extended by about 40 GeV to masses of 270 – 280 GeV. In this report event yields are presented for beam energies in the range $E_e = 27.5\text{--}35$ GeV and $E_p = 820\text{--}1100$ GeV. Also presented is the rate of background events expected from Standard Model neutral and charged current $e^\pm p$ scattering.

1 Introduction

The ZEUS [1] and H1 [2] experiments, studying deep-inelastic positron proton scattering (DIS) at HERA, recently reported an excess of events in the region of high Bjorken- x and large negative photon virtualities (Q^2)¹. This excess may result from the production of a R-parity violating squark $\tilde{q}$ or of a leptoquark LQ via the fusion of a positron and a quark (see Fig. 1),

$$e^+ p \rightarrow \tilde{q}/LQ X \rightarrow e' + h + X. \quad (1)$$

¹Preliminary results obtained recently after more than doubling the integrated luminosity did not show an increase of the excess, see [3, 4].

For the purpose of this note, $\tilde{q}$ and LQ are interchangeable. The symbol LQ will be used to denote either one of them. The data indicate for the mass of the LQ a value around $M_{LQ} = 200\text{--}230$ GeV which corresponds to a proton momentum fraction x carried by the fusing quark of $x = M_{LQ}^2/s = 0.45\text{--}0.60$, where $s = 4E_e E_p$ is the square of the total ep c.m. energy and E_e, E_p are the positron and proton beam energies. The possible existence of $\tilde{q}$ or LQ in this mass range prompted us to study the $\tilde{q}/LQ$ production rate as a function of the HERA beam energies.

The lowest order cross section for producing a scalar R-parity violating squark $\tilde{q}$ or leptoquark LQ in ep collisions can be written as (see e.g. [5])

$$\frac{d\sigma_{ep \rightarrow LQ}}{dy} = \frac{\pi}{4s} \lambda^2 q(x, Q^2) = \frac{\pi}{4M_{LQ}^2} \lambda^2 x q(x, Q^2) \quad (2)$$

where $Q^2 = -(e - e')^2$ and $y = Q^2/(xs)$ are the standard DIS variables and $xq(x, Q^2)$ is the momentum density of the quark of type q in the proton fusing with an electron or positron, generically denoted by e ; λ is a coupling constant which depends on the type of quark and lepton that form the $\tilde{q}/LQ$ state. The large value of x plus the small production cross section combined with the absence of a signal in the electron-proton data favors the assumption that the fusing quark is an up or down quark (see e.g. [6]). Since the density of anti-sea quarks is small at large x we expect that the LQ production cross section is proportional to the momentum densities of the quarks, $xu_v(x, Q^2)$, $xd_v(x, Q^2)$ or $xs(x, Q^2)$, which are steeply falling with x at high x . For this reason, a large increase of the production cross section can be obtained by a moderate increase of the HERA beam energies.

This note presents the expected increase of the $\tilde{q}/LQ$ production cross section and the $\tilde{q}/LQ$ search range for various combinations of positron and proton beam energies. Also given are the number of background events predicted by the Standard Model (SM) for neutral current (NC) and charged current (CC) $e^\pm p$ scattering. The data reported in [1, 2] were taken at the ('nominal') beam energies of $E_e = 27.5$ GeV, $E_p = 820$ GeV. The range explored in this study is $E_e = 27.5\text{--}35$ GeV and $E_p = 820\text{--}1100$ GeV. The maximum values of $E_e = 35$ GeV and $E_p = 1100$ GeV should be considered as extreme upper limits for HERA.

2 Choice of parameters

The scalar LQ (or $\tilde{q}$) production cross sections were calculated from eq. (2) for the processes $e^+d \rightarrow LQ$, $e^+u \rightarrow LQ$ and $e^+s \rightarrow LQ$. The quark momentum densities were taken from the parton distribution set MRSA [8]. In order to fix the production rate of LQ events the LQ cross section was integrated over $y > 0.25$ and normalized such that the number of LQ candidates accepted in e^+p NC scattering as reported in ref. [1, 2, 3, 4] is approximately reproduced. Taking $M_{LQ} = 215$ GeV and assuming 10 LQ events for an integrated luminosity of $L = 50 \text{ pb}^{-1}$ at the nominal beam energies $E_e (E_p) = 27.5 (820)$ GeV yields for the coupling constant λ the values 0.045, 0.021, 0.7 for the e^+d ,

e^+u and e^+s states, respectively. For simplicity, we assume that the acceptance for LQ candidates does not depend on whether a NC or a CC event sample is selected.

The background from the standard NC processes, $e^\pm p \rightarrow e^\pm X$, was estimated by calculating the differential cross sections

$$\frac{d^2\sigma_{NC}^\pm}{dxdy} = \frac{2\pi\alpha^2 s Y_+}{Q^4} \left\{ F_{2,NC} - \frac{y^2}{Y_+} F_{L,NC} \mp \frac{Y_-}{Y_+} x F_{3,NC} \right\}, \quad (3)$$

taking into account the contributions from γZ interference and Z exchange [9]. Likewise the cross sections

$$\frac{d^2\sigma_{CC}^\pm}{dxdy} = \frac{G_F^2 s Y_+}{8\pi} \frac{M_W^4}{(Q^2 + M_W^2)^2} \left\{ F_{2,CC}^\pm - \frac{y^2}{Y_+} F_{L,CC}^\pm \mp \frac{Y_-}{Y_+} x F_{3,CC}^\pm \right\} \quad (4)$$

were calculated for the CC processes $e^+p \rightarrow \bar{\nu}X$ and $e^-p \rightarrow \nu X$. In eqs. (3) and (4) α is the fine structure constant, G_F the Fermi constant, M_W the mass of the W and $Y_\pm = 1 \pm (1-y)^2$. The structure functions F_2 and xF_3 were calculated in leading order with the parton distributions of the proton taken from MRSA. The contribution from F_L in this region of high x is expected to be small and was neglected. The cross sections were integrated over $y > 0.25$ and over the band $x = M_{LQ}^2/s \pm 0.08$ which corresponds roughly to ± 2 standard deviations for the LQ mass measurement of [1, 2]. The number of background events was obtained by normalizing the e^+p NC cross section to a yield of 0.675 accepted events in the bin $0.55 < x < 0.65$, $y > 0.25$ and $L = 20 \text{ pb}^{-1}$ [1]. In this way the acceptance quoted in [1] for e^+p NC scattering is approximately taken into account. The same acceptance was assumed for e^-p NC and $e^\pm p$ CC scattering.

3 Signal and background for $M_{LQ} = 200 - 230 \text{ GeV}$

Tables 1 and 2 give the estimated event yields (N_{LQ}^i) for $e^+q_i \rightarrow LQ$, $q_i = d, u, s$ with an integrated luminosity of $L = 50 \text{ pb}^{-1}$ for various combinations of $E_e = 27.5\text{--}35 \text{ GeV}$, $E_p = 820\text{--}1100 \text{ GeV}$ and $M_{LQ} = 200\text{--}230 \text{ GeV}$. Also given are the expected number of e^+p and e^-p NC and CC background events ($N_{NC,CC}^+$, $N_{NC,CC}^-$). It is seen that for e^+p scattering the CC background is about a factor of five lower than the NC background whereas in e^-p scattering the CC and NC contributions are roughly equal in size. The total background in e^+p scattering is about a factor of four lower than in e^-p .

Tables 3 and 4 present the gain in LQ production, defined as

$$G(E_e, E_p, M_{LQ}) = \frac{N_{LQ}(E_e, E_p, M_{LQ})}{N_{LQ}(E_e^0, E_p^0, M_{LQ})}$$

where E_e^0 (E_p^0) = 27.5 (820) GeV denote the nominal beam energies. Also listed are the signal over background ratios

$$S_{CC,NC}^+ = \frac{N_{LQ}}{N_{CC,NC}^+}$$

for e^+p scattering.

In Figs. 2 and 3 we show the gain in LQ events with respect to production at nominal beam energies for $e^+d \rightarrow LQ$ and $e^+u \rightarrow LQ$, respectively. The gain is larger for larger leptoquark masses mainly because the shift towards smaller x_{LQ} is larger when M_{LQ} increases, see Tables 1 and 2. Comparing Figs. 2 and 3 it is seen that a larger gain is obtained for the e^+d as compared to the e^+u case. This is so because the d_v distribution is considerably steeper at high x than the u_v distribution as illustrated in Fig. 5a. For $e^+s \rightarrow LQ$ production the gain is particularly large (see Fig. 4) because the strange quark distribution is a very steep function of x at large x , as shown in Fig. 5b. However, the strange quark distribution in the proton is not very well known. This is illustrated in Fig. 5b, where the distribution from MRSA is compared with those from CTEQ3M [10] and GRV [11]: the gain in LQ production from s quarks is about a factor of 1.5 lower if CTEQ3M instead of MRSA parton distributions are used.

Figures 6–9 show the e^+ and e^- NC and CC ‘reduced’ cross sections defined as the terms inside the curly brackets in eqs. (3) and (4) for $s = 9 \times 10^4 \text{ GeV}^2$ (corresponding to the nominal beam energies) and $s = 15 \times 10^4 \text{ GeV}^2$ ($E_e = 35 \text{ GeV}$, $E_p = 1100 \text{ GeV}$). For e^+p scattering, the reduced NC and CC cross sections at fixed Q^2 above the masses squared of W , Z increase with s while for e^-p scattering they decrease.

4 Search range for squarks/leptoquarks at HERA

The cross sections for production of squarks/leptoquarks in ep collisions by ed , eu and es fusion are shown in Figs. 10, 11, 12 as a function of M_{LQ} between 200 and 300 GeV for different combinations of beam energies. They were calculated from Eq. 2, using the MRSA parton distribution set, and integrated over $y > 0.25$. As before, we assumed $\lambda_{ed} = 0.045$, $\lambda_{eu} = 0.021$, $\lambda_{es} = 0.7$. Assuming that a $\tilde{q}/LQ$ signal can be established with 10 events, a minimum cross section of 50 fb is required for an integrated luminosity of 200 pb^{-1} . For the nominal beam energies ($E_e = 27.5 \text{ GeV}$, $E_p = 820 \text{ GeV}$) this condition limits the search range to masses of 225 - 235 GeV. By raising the beam energies to (32.5 GeV, 1000 GeV) the search range is extended by about 40 GeV to 270 - 280 GeV. If the beam energies are raised even higher to (35 GeV, 1100 GeV) the search range increases to $M_{LQ} = 290 - 300 \text{ GeV}$. Of course, the search range is larger (smaller) for higher (lower) values of λ .

In this study a minimum of 10 signal events were required using the LO cross section for LQ production as given by Eq. 2. QCD corrections from gluon radiation, gluon splitting and vertex corrections increase the cross section [12, 13]. However, the requirement of a minimum of 10 signal events leads basically to the same conclusions for the search range as before.

5 Conclusions

The production rate of squarks/leptoquarks with a mass around 200 GeV can be increased substantially by raising the beam energies of HERA. The number of LQ events increases

by a factor of 1.5 for $E_e, E_p = 27.5$ GeV, 900 GeV	
2.1	27.5 GeV, 1000 GeV
2.0	30.0 GeV, 900 GeV
2.7	30.0 GeV, 1000 GeV
3.3	32.5 GeV, 1000 GeV.

These numbers were obtained for $M_{LQ} = 215$ GeV and represent averages over e^+u and e^+d production. For e^+s leptoquark production the gains are even larger, see Fig. 4.

By raising the beam energies from $E_e = 27.5$ GeV, $E_p = 820$ GeV to $E_e = 32.5$ GeV, $E_p = 1000$ GeV (35 GeV, 1100 GeV) the search range for squarks/leptoquarks can be extended by about 40 (75) GeV to $M_{LQ} = 270 - 280$ (290 - 300) GeV.

6 Acknowledgements

We thank Dr. U. Katz for useful comments.

References

- [1] ZEUS Collaboration, J. Breitweg et al., DESY 97-025.
- [2] H1 Collaboration, S. Adloff et al., DESY 97-024.
- [3] A.T. Doyle, *Highlights and Open Questions from ZEUS*, Proc. DIS98 Workshop, Brussels, 1998.
- [4] T. Greenshaw, *Results from H1*, Proc. DIS98 Workshop, Brussels, 1998.
- [5] W. Buchmüller, R. Rückl and D. Wyler, Phys. Lett. B 191 (1987) 442.
- [6] K.S. Babu, C. Kolda, J. March-Russell and F. Wilczek, Phys. Lett. B402 (1997) 367.
- [7] G. Altarelli, J. Ellis, G.F. Giudice, S. Lola and M.L. Mangano, Nucl. Phys. B506 (1997) 3.
- [8] A.D Martin, W.J. Stirling and R.G. Roberts, Phys. Rev. D50 (1994) 6734.

- [9] G. Ingelman and R. Rückl, Z. Phys. C44 (1989) 291;
J. Blümlein et al., Z. Phys. C45 (1990) 501.
- [10] CTEQ Collaboration, J. Botts et al., Phys. Lett. B304 (1993) 159.
- [11] M. Glück, E. Reya and A. Vogt, Z. Phys. C67 (1995) 433.
- [12] T. Plehn, H. Spiesberger, M. Spira and P. Zerwas, Z. Phys. C74 (1997) 611.
- [13] Z. Kunszt and W.J. Stirling, Z. Phys. C75 (1997) 453.

E_e (GeV)	E_p (GeV)	M_{LQ} (GeV)	x_{LQ}	$e^+q_i \rightarrow LQ$			$ep \rightarrow e(\nu)X$			
				N_{LQ}^d	N_{LQ}^u	N_{LQ}^s	N_{NC}^+	N_{CC}^+	N_{NC}^-	N_{CC}^-
27.5	820	200	0.44	10.0	10.0	10.0	20.1	4.4	39.0	53.7
27.5	820	210	0.49	10.0	10.0	10.0	11.8	2.5	23.5	33.9
27.5	820	220	0.54	10.0	10.0	10.0	6.7	1.3	13.5	20.3
27.5	820	230	0.59	10.0	10.0	10.0	3.6	0.6	7.3	11.4
27.5	900	200	0.40	13.7	12.7	18.8	28.7	7.0	55.2	75.1
27.5	900	210	0.45	14.8	13.5	21.3	17.8	4.1	35.1	50.1
27.5	900	220	0.49	16.1	14.6	24.7	10.7	2.3	21.5	32.2
27.5	900	230	0.53	18.0	16.0	30.0	6.2	1.3	12.7	19.7
27.5	1000	200	0.36	18.5	15.9	34.6	41.1	11.3	78.1	104.6
27.5	1000	210	0.40	21.2	17.8	43.5	26.6	7.0	51.9	73.2
27.5	1000	220	0.44	24.9	20.4	57.6	16.9	4.2	33.8	49.9
27.5	1000	230	0.48	30.5	24.3	82.0	10.5	2.5	21.4	32.8
27.5	1100	200	0.33	23.1	18.6	55.4	55.1	16.7	103.3	136.3
27.5	1100	210	0.36	27.7	21.6	75.3	36.8	10.7	71.0	98.7
27.5	1100	220	0.40	34.3	26.0	109.3	24.3	6.7	48.1	70.0
27.5	1100	230	0.44	44.9	32.7	173.4	15.8	4.1	31.9	48.4
30.0	820	200	0.41	13.5	12.5	18.1	28.1	6.8	54.1	73.6
30.0	820	210	0.45	14.4	13.3	20.3	17.4	4.0	34.2	48.9
30.0	820	220	0.49	15.7	14.2	23.4	10.4	2.3	20.9	31.3
30.0	820	230	0.54	17.4	15.6	28.1	6.0	1.2	12.3	19.1
30.0	900	200	0.37	17.6	15.3	31.3	38.7	10.4	73.7	99.0
30.0	900	210	0.41	20.0	17.0	38.8	24.9	6.4	48.7	68.8
30.0	900	220	0.45	23.2	19.3	50.3	15.7	3.8	31.4	46.4
30.0	900	230	0.49	28.1	22.7	69.8	9.7	2.2	19.6	30.2
30.0	1000	200	0.33	22.7	18.4	53.3	53.8	16.2	100.9	133.3
30.0	1000	210	0.37	27.1	21.3	72.0	35.8	10.3	69.2	96.3
30.0	1000	220	0.40	33.5	25.5	103.8	23.6	6.5	46.7	68.1
30.0	1000	230	0.44	43.5	31.9	163.3	15.3	4.0	30.9	46.9
30.0	1100	200	0.30	27.5	21.0	80.7	70.8	23.4	130.7	169.7
30.0	1100	210	0.33	34.0	25.1	116.2	48.3	15.4	91.9	126.1
30.0	1100	220	0.37	43.9	31.2	180.6	32.8	10.0	64.1	92.1
30.0	1100	230	0.40	60.2	40.8	310.4	22.0	6.4	44.0	66.0

Table 1: The expected number of events N_{LQ}^i for the squark/leptoquark production processes $e^+q_i \rightarrow LQ$, $q_i = d, u, s$. The number of background events from $e^\pm p$ NC and CC scattering are listed in the columns $N_{NC}^\pm$ and $N_{CC}^\pm$. The event yields are given in the bin $x_{LQ} \pm 0.08$, $y > 0.25$ for an integrated luminosity of $L = 50 \text{ pb}^{-1}$.

E_e (GeV)	E_p (GeV)	M_{LQ} (GeV)	x_{LQ}	$e^+q_i \rightarrow LQ$			$ep \rightarrow e(\nu)X$			
				N_{LQ}^d	N_{LQ}^u	N_{LQ}^s	N_{NC}^+	N_{CC}^+	N_{NC}^-	N_{CC}^-
32.5	820	200	0.38	17.0	14.9	29.2	37.1	9.8	70.7	95.2
32.5	820	210	0.41	19.1	16.5	35.6	23.7	6.0	46.5	65.8
32.5	820	220	0.45	22.1	18.6	45.6	14.9	3.6	29.8	44.1
32.5	820	230	0.50	26.4	21.7	62.1	9.1	2.0	18.5	28.5
32.5	900	200	0.34	21.5	17.7	47.3	49.8	14.6	93.9	124.5
32.5	900	210	0.38	25.3	20.3	62.6	33.0	9.2	63.8	89.2
32.5	900	220	0.41	30.9	24.0	88.2	21.5	5.7	42.7	62.4
32.5	900	230	0.45	39.5	29.6	135.1	13.8	3.5	27.9	42.5
32.5	1000	200	0.31	26.7	20.6	75.8	67.9	22.1	125.6	163.5
32.5	1000	210	0.34	32.9	24.5	108.1	46.1	14.4	88.0	121.0
32.5	1000	220	0.37	42.2	30.3	166.2	31.2	9.3	61.0	88.0
32.5	1000	230	0.41	57.3	39.3	281.9	20.8	5.9	41.7	62.7
32.5	1100	200	0.28	31.5	23.0	109.9	88.1	31.3	159.9	204.5
32.5	1100	210	0.31	40.0	28.1	165.5	61.0	21.0	114.4	154.8
32.5	1100	220	0.34	53.3	35.9	271.0	42.2	13.9	81.4	115.7
32.5	1100	230	0.37	75.6	48.3	494.8	29.0	9.2	57.4	85.2
35.0	820	200	0.35	20.6	17.1	43.1	47.0	13.5	88.8	118.2
35.0	820	210	0.38	24.0	19.5	56.2	30.9	8.5	60.0	84.1
35.0	820	220	0.42	29.0	22.9	77.8	20.0	5.2	39.8	58.4
35.0	820	230	0.46	36.6	27.9	116.6	12.7	3.1	25.8	39.4
35.0	900	200	0.32	25.1	19.7	66.4	62.1	19.6	115.5	151.3
35.0	900	210	0.35	30.6	23.3	92.8	41.9	12.7	80.3	111.0
35.0	900	220	0.38	38.7	28.4	139.2	28.0	8.1	55.2	79.9
35.0	900	230	0.42	51.8	36.4	229.8	18.5	5.1	37.3	56.2
35.0	1000	200	0.29	30.4	22.5	101.6	83.2	29.0	151.7	194.9
35.0	1000	210	0.32	38.4	27.4	151.3	57.4	19.3	108.1	146.8
35.0	1000	220	0.35	50.8	34.7	244.5	39.5	12.8	76.6	109.1
35.0	1000	230	0.38	71.4	46.3	439.8	27.0	8.3	53.7	79.8
35.0	1100	200	0.26	35.1	24.7	142.2	107.1	40.5	191.0	240.4
35.0	1100	210	0.29	45.6	30.8	222.1	74.9	27.6	138.4	184.7
35.0	1100	220	0.31	62.3	40.0	378.9	52.5	18.7	100.1	140.5
35.0	1100	230	0.34	90.9	55.2	725.7	36.8	12.5	72.0	105.5

Table 2: Continuation of Table 1.

E_e (GeV)	E_p (GeV)	M_{LQ} (GeV)	x_{LQ}	$e^+d \rightarrow LQ$			$e^+u \rightarrow LQ$			$e^+s \rightarrow LQ$		
				G	S_{NC}^+	S_{CC}^+	G	S_{NC}^+	S_{CC}^+	G	S_{NC}^+	S_{CC}^+
27.5	820	200	0.44	1.0	0.5	2.3	1.0	0.5	2.3	1.0	0.5	2.3
27.5	820	210	0.49	1.0	0.8	4.1	1.0	0.8	4.1	1.0	0.8	4.1
27.5	820	220	0.54	1.0	1.5	7.7	1.0	1.5	7.7	1.0	1.5	7.7
27.5	820	230	0.59	1.0	2.8	15.5	1.0	2.8	15.5	1.0	2.8	15.5
27.5	900	200	0.40	1.4	0.5	2.0	1.3	0.4	1.8	1.9	0.7	2.7
27.5	900	210	0.45	1.5	0.8	3.6	1.4	0.8	3.3	2.1	1.2	5.1
27.5	900	220	0.49	1.6	1.5	6.9	1.5	1.4	6.2	2.5	2.3	10.5
27.5	900	230	0.53	1.8	2.9	14.1	1.6	2.6	12.6	3.0	4.8	23.5
27.5	1000	200	0.36	1.9	0.5	1.6	1.6	0.4	1.4	3.5	0.8	3.1
27.5	1000	210	0.40	2.1	0.8	3.0	1.8	0.7	2.5	4.4	1.6	6.2
27.5	1000	220	0.44	2.5	1.5	5.9	2.0	1.2	4.8	5.8	3.4	13.7
27.5	1000	230	0.48	3.1	2.9	12.4	2.4	2.3	9.8	8.2	7.8	33.3
27.5	1100	200	0.33	2.3	0.4	1.4	1.9	0.3	1.1	5.5	1.0	3.3
27.5	1100	210	0.36	2.8	0.8	2.6	2.2	0.6	2.0	7.5	2.0	7.0
27.5	1100	220	0.40	3.4	1.4	5.1	2.6	1.1	3.9	10.9	4.5	16.2
27.5	1100	230	0.44	4.5	2.8	10.8	3.3	2.1	7.9	17.3	11.0	41.8
30.0	820	200	0.41	1.3	0.5	2.0	1.3	0.4	1.8	1.8	0.6	2.7
30.0	820	210	0.45	1.4	0.8	3.6	1.3	0.8	3.3	2.0	1.2	5.1
30.0	820	220	0.49	1.6	1.5	6.9	1.4	1.4	6.3	2.3	2.2	10.4
30.0	820	230	0.54	1.7	2.9	14.2	1.6	2.6	12.7	2.8	4.7	23.0
30.0	900	200	0.37	1.8	0.5	1.7	1.5	0.4	1.5	3.1	0.8	3.0
30.0	900	210	0.41	2.0	0.8	3.1	1.7	0.7	2.7	3.9	1.6	6.1
30.0	900	220	0.45	2.3	1.5	6.1	1.9	1.2	5.1	5.0	3.2	13.2
30.0	900	230	0.49	2.8	2.9	12.7	2.3	2.4	10.3	7.0	7.2	31.6
30.0	1000	200	0.33	2.3	0.4	1.4	1.8	0.3	1.1	5.3	1.0	3.3
30.0	1000	210	0.37	2.7	0.8	2.6	2.1	0.6	2.1	7.2	2.0	7.0
30.0	1000	220	0.40	3.3	1.4	5.2	2.6	1.1	3.9	10.4	4.4	16.0
30.0	1000	230	0.44	4.4	2.8	11.0	3.2	2.1	8.0	16.3	10.7	41.1
30.0	1100	200	0.30	2.7	0.4	1.2	2.1	0.3	0.9	8.1	1.1	3.5
30.0	1100	210	0.33	3.4	0.7	2.2	2.5	0.5	1.6	11.6	2.4	7.6
30.0	1100	220	0.37	4.4	1.3	4.4	3.1	1.0	3.1	18.1	5.5	18.1
30.0	1100	230	0.40	6.0	2.7	9.4	4.1	1.9	6.4	31.0	14.1	48.7

Table 3: The gain $G = N_{LQ}(E_e, E_p, M_{LQ})/N_{LQ}(27.5, 820, M_{LQ})$ and the signal to background ratios $S_{NC,CC}^+ = N_{LQ}/N_{NC,CC}^+$ for the processes $e^+d \rightarrow LQ$, $e^+u \rightarrow LQ$ and $e^+s \rightarrow LQ$.

E_e (GeV)	E_p (GeV)	M_{LQ} (GeV)	x_{LQ}	$e^+d \rightarrow LQ$			$e^+u \rightarrow LQ$			$e^+s \rightarrow LQ$		
				G	S_{NC}^+	S_{CC}^+	G	S_{NC}^+	S_{CC}^+	G	S_{NC}^+	S_{CC}^+
32.5	820	200	0.38	1.7	0.5	1.7	1.5	0.4	1.5	2.9	0.8	3.0
32.5	820	210	0.41	1.9	0.8	3.2	1.6	0.7	2.7	3.6	1.5	5.9
32.5	820	220	0.45	2.2	1.5	6.2	1.9	1.2	5.2	4.6	3.1	12.8
32.5	820	230	0.50	2.6	2.9	12.9	2.2	2.4	10.6	6.2	6.8	30.4
32.5	900	200	0.34	2.1	0.4	1.5	1.8	0.4	1.2	4.7	0.9	3.2
32.5	900	210	0.38	2.5	0.8	2.7	2.0	0.6	2.2	6.3	1.9	6.8
32.5	900	220	0.41	3.1	1.4	5.4	2.4	1.1	4.2	8.8	4.1	15.4
32.5	900	230	0.45	4.0	2.9	11.4	3.0	2.1	8.5	13.5	9.8	38.9
32.5	1000	200	0.31	2.7	0.4	1.2	2.1	0.3	0.9	7.6	1.1	3.4
32.5	1000	210	0.34	3.3	0.7	2.3	2.5	0.5	1.7	10.8	2.3	7.5
32.5	1000	220	0.37	4.2	1.4	4.5	3.0	1.0	3.2	16.6	5.3	17.8
32.5	1000	230	0.41	5.7	2.8	9.7	3.9	1.9	6.6	28.2	13.5	47.6
32.5	1100	200	0.28	3.1	0.4	1.0	2.3	0.3	0.7	11.0	1.2	3.5
32.5	1100	210	0.31	4.0	0.7	1.9	2.8	0.5	1.3	16.6	2.7	7.9
32.5	1100	220	0.34	5.3	1.3	3.8	3.6	0.9	2.6	27.1	6.4	19.4
32.5	1100	230	0.37	7.6	2.6	8.3	4.8	1.7	5.3	49.5	17.1	54.0
35.0	820	200	0.35	2.1	0.4	1.5	1.7	0.4	1.3	4.3	0.9	3.2
35.0	820	210	0.38	2.4	0.8	2.8	2.0	0.6	2.3	5.6	1.8	6.6
35.0	820	220	0.42	2.9	1.4	5.5	2.3	1.1	4.4	7.8	3.9	14.9
35.0	820	230	0.46	3.7	2.9	11.7	2.8	2.2	8.9	11.7	9.2	37.2
35.0	900	200	0.32	2.5	0.4	1.3	2.0	0.3	1.0	6.6	1.1	3.4
35.0	900	210	0.35	3.1	0.7	2.4	2.3	0.6	1.8	9.3	2.2	7.3
35.0	900	220	0.38	3.9	1.4	4.8	2.8	1.0	3.5	13.9	5.0	17.2
35.0	900	230	0.42	5.2	2.8	10.2	3.6	2.0	7.2	23.0	12.4	45.1
35.0	1000	200	0.29	3.0	0.4	1.0	2.2	0.3	0.8	10.2	1.2	3.5
35.0	1000	210	0.32	3.8	0.7	2.0	2.7	0.5	1.4	15.1	2.6	7.8
35.0	1000	220	0.35	5.1	1.3	4.0	3.5	0.9	2.7	24.4	6.2	19.1
35.0	1000	230	0.38	7.1	2.6	8.6	4.6	1.7	5.6	44.0	16.3	52.7
35.0	1100	200	0.26	3.5	0.3	0.9	2.5	0.2	0.6	14.2	1.3	3.5
35.0	1100	210	0.29	4.6	0.6	1.7	3.1	0.4	1.1	22.2	3.0	8.1
35.0	1100	220	0.31	6.2	1.2	3.3	4.0	0.8	2.1	37.9	7.2	20.3
35.0	1100	230	0.34	9.1	2.5	7.2	5.5	1.5	4.4	72.6	19.7	57.9

Table 4: Continuation of table 3.




Figure 1: Diagram for leptoquark production by ep scattering.




Figure 2: The gain $G = N_{LQ}(E_e, E_p)/N_{LQ}(27.5, 820)$ in squark/leptoquark production by e^+d scattering as function of the positron energy E_e for various proton energies E_p and assumed squark/leptoquark masses of (a) $M_{LQ} = 200 \text{ GeV}$; (b) 210 GeV ; (c) 220 GeV and (d) 230 GeV .




Figure 3: The gain $G = N_{LQ}(E_e, E_p)/N_{LQ}(27.5, 820)$ in squark/leptoquark production by e^+u scattering as function of the positron energy E_e for various proton energies E_p and assumed squark/leptoquark masses of (a) $M_{LQ} = 200 \text{ GeV}$; (b) 210 GeV ; (c) 220 GeV and (d) 230 GeV .




Figure 4: The gain $G = N_{LQ}(E_e, E_p)/N_{LQ}(27.5, 820)$ in squark/leptoquark production by e^+s scattering as function of the positron energy E_e for various proton energies E_p and assumed squark/leptoquark masses of (a) $M_{LQ} = 200 \text{ GeV}$; (b) 210 GeV ; (c) 220 GeV and (d) 230 GeV .




Figure 5: (a) The down (solid curve) and up (dashed curve) valence distributions and (b) the strange quark distribution versus x at a fixed value of $Q^2 = 3 \times 10^4 \text{ GeV}^2$. The distributions are taken from the parton distribution set MRSA and are normalized to unity at $x = 0.6$. Also shown are the strange quark distributions from the sets CTEQ3M and GRV.




Figure 6: The reduced cross section (see text) for the NC process $e^+p \rightarrow e^+X$.




Figure 7: The reduced cross section (see text) for the CC process $e^+p \rightarrow \bar{\nu}X$.




Figure 8: The reduced cross section (see text) for the NC process $e^-p \rightarrow e^-X$.




Figure 9: The reduced cross section (see text) for the CC process $e^-p \rightarrow \nu X$.




Figure 10: The cross section $\sigma(ep \rightarrow LQ X)$ for squark/leptoquark production by ed fusion for $\lambda_{ed} = 0.045$; see text.




Figure 11: The cross section $\sigma(ep \rightarrow LQ X)$ for squark/leptoquark production by eu fusion for $\lambda_{eu} = 0.021$; see text.




Figure 12: The cross section $\sigma(ep \rightarrow LQ X)$ for squark/leptoquark production by es fusion for $\lambda_{es} = 0.7$; see text.